

GENERAL INSTRUCTIONS

1. **Bring an adult driver.** Your driver must be 18 years of age or over and they must stay in the Endoscopy Center. You will be given a sedative for your test, and you are not to drive until the next day. Therefore, one person must stay and drive you home. Your procedure will be rescheduled if your driver cannot stay in the Center.
2. **If you need to reschedule** your procedure, please call your physician's office at least 3-4 days in advance so that we may schedule another patient in your appointment time.
3. **NOTIFY US IN ADVANCE:**
 - If you are taking a **blood thinner** prescribed by a physician (such as Coumadin, Plavix, Effient, Eliquis or Ticlid). Your gastroenterologist will need to determine if these medication dosages require adjustment prior to the procedure. Failure to properly adjust these medications may result in cancellation of your procedure. You may continue to take Aspirin as usual.
 - If you are having **chest pain or undergoing cardiac testing**. (IMPORTANT: Please notify us before you begin to prep for your procedure.)
 - If you have ever had **excessive bleeding after an operation or problems with anesthesia**.
 - If there is a possibility you are **pregnant**.
4. **DO NOT:**
 - Take **iron, diet pills** (Phentermine, Adipex or Qsymia) or **herbal medications** for seven (7) days prior to having your procedure. You may take vitamins.
 - Drink **alcohol** for 24 hours before or after your procedure.
 - Use items such as **gum, hard candy, breath mints, smokeless tobacco, or illicit drugs** such as marijuana on the day of your procedure.
5. If you have **dentures**, you may be required to remove them for the procedure.
6. **Leave your valuables at home.** If you have **body piercings**, you will be asked to remove them prior to the procedure for your safety.
7. If you are a **diabetic**, do not take your diabetic medication the day of your procedure. Bring your medication with you so you can take it after your procedure.
8. If you have **asthma**, please bring your inhaler with you.
9. Take your usual **prescribed medications** with no more than two (2) tablespoons / one (1) ounce of water up to three (3) hours prior to your appointment time.



GASTROENTEROLOGY ASSOCIATES
OF NORTH MISSISSIPPI

powered by



GLP-1 DRUGS *** Hold as Instructed Prior to Procedure ***

The following weight loss / diabetic drugs should be held as instructed prior to procedure:

Liraglutide: Victoza (weekly) and Saxenda (daily)

Dulaglutide: Trulicity (weekly)

Exenatide: Bydureon (weekly) and Byetta (daily)

Semaglutide: Ozempic (weekly), Wegovy (weekly) and Rybelsus (daily)

Tirzepatide: Mounjaro (weekly) and Zepbound (weekly)

Lixisenatide: Adlyxin (daily)

The medication will need to be **stopped seven (7) days prior** to your procedure if you are on a weekly dose or **the day of** your procedure if you are on a daily dose.

For any questions or to reschedule your appointment, please call our office at 662-234-9888. We apologize for any inconvenience this may cause.

Please call us at 662-234-9888 if you have any questions or if you are having difficulty with the preparation. We are available 24 hours a day, 7 days a week.

Clenpiq Prep

5 - 7 DAYS BEFORE THE PROCEDURE - Have your prescription filled. Please follow the prep instructions listed below, not the instructions on the prep box. Do NOT refrigerate the prep.

5 DAYS BEFORE THE PROCEDURE - If you routinely have less than two (2) bowel movements per week then you will also need to purchase an 8.3oz bottle of Miralax. Begin taking a dose of Miralax twice a day **5 days prior** to your procedure. Take the last dose of Miralax in the morning **the day before** your procedure.

3 DAYS BEFORE THE PROCEDURE - AVOID raw fruits and vegetables, whole wheat/multigrain products, beans, nuts, popcorn, and seeds.

THE DAY BEFORE THE PROCEDURE - You may only have **clear liquids the entire day** such as water, coffee, tea, broth, sodas, apple juice, Gatorade, Kool-Aid, Jell-O, popsicles, and hard candy. You may not have any solid food. **It is important to drink extra clear liquids before your prep to maintain adequate hydration. Do not drink liquids that are not clear such as milk. Do not drink liquids that are red or purple in color.**

1 DAY BEFORE THE PROCEDURE – BETWEEN 3PM & 6PM – START YOUR PREP

STEP 1:

- Take the first dose (5.4 oz bottle) of CLENPIQ.
- Drink the entire contents. A straw may help.

STEP 2:

- Follow with at least FIVE (5) 8oz cups of clear liquid, taken at your own pace before going to bed.
- **Make a conscious effort to drink as much clear liquids as you can before, during and after the preparation.**



NOTE:

- If you experience nausea - slow down your prep intake and walk around until nausea is relieved.
- Use baby wipes and/or Balneol cleansing cream or lotion during the prep to reduce irritation.

DAY OF PROCEDURE - 4 HOURS BEFORE YOU LEAVE HOME FOR APPOINTMENT

STEP 1:

- Take the second dose (5.4oz bottle) of CLENPIQ.
- Drink the entire contents. A straw may help.

STEP 2:

- Follow with at least THREE (3) 8oz cups of clear liquid within the next hour. You may drink clear liquids until 3 hours before your appointment time.
- You may take your prescription medications (except those listed in the booklet) with water up to three (3) hours before your appointment arrival time.
- Do not use smokeless tobacco on the day of your procedure.
- Starting three (3) hours before your appointment arrival time **absolutely nothing by mouth**, including gum, hard candy, breath mints.



Please call us at 662-234-9888 if you have any questions or if you are having difficulty with the preparation. We have someone available 24 hours a day, 7 days a week that can assist you.

Diabetic Prep

MEDICATIONS:

- If you take oral medications for your diabetes, do NOT take your diabetes pills on the day prior to your procedure (prep day) or on the day of your procedure (until your procedure is complete and you resume a normal diet).
- If you take non-insulin injectable medications (Byetta, Bydureon, Victoza, Trulicity, Tanzeum, Symlin), do not take the medication if your injection day is the day prior to your procedure (prep day). You can resume it after the procedure is complete.
- If you take insulin, reduce your dose by 50% (half) of your normal dose on the day prior to your procedure and the day of your procedure. You can resume your normal dose after the procedure when you resume a normal diet.
- If you use an insulin pump, contact your diabetes doctor for instruction. Generally, most patients will need to reduce their basal rate to 80% of the normal rate.

DIET:

- You should aim for 45-60 grams of carbs with each meal. Although you may ordinarily avoid sugars, they are ok to take when you are prepping for your colonoscopy as these will be the main source of calories when you are on a clear liquid diet.
- For snacks, try to include other fluids that contain electrolytes such as broth.
- Examples of clear liquids with carb amounts are listed below. Note, if you experience low blood sugar (<60) during the 2 hours before your procedure, you can take 3 teaspoons of soda or clear juice to raise your sugar. Do NOT take solid foods or non-clear liquids, as your procedure will be cancelled.

FLUID	AMOUNT	CARBS (grams)
Juice	½ cup	15g
Jello	½ cup	15g
Soda	1 can	40g
Popsicle	1 double stick	15g
Gatorade / Powerade	½ cup	15g
Broth (veg / chicken / beef)	Any	0g
Clear Protein Drinks	Any	0g